

Galactic Bulge Observations

Thomas L. Roellig, Kin-Wing Chan

The mid-infrared spectrometer (MIRS) that flew on the joint NASA/Japanese Space Agency Infrared Telescope in Space (IRTS) mission in 1995 yielded a wealth of data that are presently being analyzed and reported on. Here we present the results of MIRS studies of the Galactic Bulge region.

The MIRS operated over wavelengths ranging from 4.6 to 11.7 microns with a spectral resolution of about 0.23 to 0.36 microns. This is an ideal spectral region and resolution for investigations of infrared emissions from solid material, since it covers many of the diagnostic stretching and bending modes of candidate interstellar material. The IRTS/MIRS telescope/instrument combination achieved sensitivities that were orders of magnitude better than those of any other instrumentation previously used for diffuse mid-infrared emission studies.

Analysis of the MIRS results was performed for four locations in the Galactic Bulge region with galactic longitude and latitude coordinates of 8.7° , 2.9° ; 8.7° , 4.0° ; 8.7° , 4.7° ; and 8.7° , 5.7° . The MIRS data were averaged over effective beam sizes of 8×20 arcminutes and corrected for interstellar extinction based on results from the Cosmic Background Explorer mission. Analysis of the data from the Infrared Astronomical Satellite (IRAS) mission indicates that the MIRS diffuse emission measurements had at most a 10% contribution from IRAS 12-micron point-sources.

Below galactic latitudes of 4.0° in the Galactic Bulge, the MIRS spectra are very similar to those from M and K giant stars. The unidentified infrared bands at 6.2, 7.7, 8.6, and 11.3 microns were also detected; it is likely they originate from emission from the diffuse interstellar medium in the bulge. Above galactic latitudes of 4.0° , the MIRS spectra are similar to the spectra of evolved stars with the high mass-loss rates seen by IRAS. One likely interpretation is that this emission arises predominantly from a large number of low-luminosity stars that were not detected by IRAS. The age of such low-luminosity stars would have to be at least 12 billion years, and the existence of such a large number of evolved stars with high mass-loss rates in the bulge would have a significant effect on our understanding of the stellar content and evolution of the Galactic Bulge. Furthermore, since the characteristics of the Galactic Bulge region are similar in many ways to those of elliptical galaxies, these results may shed light on the origin of the excess mid-infrared emission that has been observed on some elliptical galaxies.

Point of Contact: T. Roellig/K-W Chan

(650) 604-6426/3118

troellig@mail.arc.nasa.gov

kwc@ssa1.arc.nasa.gov

Photochemistry of Interstellar and Cometary Ice Analogs

Scott Sandford, Max Bernstein, Louis Allamandola, Robert Walker

Ames Research Center has made major contributions to the understanding of the composition and properties of interstellar and cometary ices. This is done, in part, by studying how laboratory ice analogs are modified by various kinds of chemical processing, for example, by ultraviolet (UV) irradiation. These laboratory studies, coupled with telescopic observations, show that the main components of cosmic ices are simple molecules like water (H_2O), methanol (CH_3OH), carbon monoxide (CO), carbon

dioxide (CO_2), and ammonia (NH_3). Ultraviolet irradiation of such ices produces "hot" hydrogen (H), carbon (C), oxygen (O), and nitrogen (N) atoms and radicals like formyl radical (HCO) which ultimately combine with other species in the ice to make new, often more complex, organic molecules. Interstellar materials made in this manner may ultimately seed planetary systems that form within dense clouds with molecules crucial to the origin of life.